

# GONAPAN-DF

Diloxanide Furoate B P  
Metronidazole B P

گوناپین۔ ڈی ایف

دیکھا سانا ڈی فورو ایٹ بی۔ پی۔  
میٹرونیڈازول بی۔ پی۔

**GONAPAN-DF** is a new approach in the management of amoebiasis offering definite therapeutic benefits over the previously available regimens. It is a combination of Diloxanide Furoate, an outstanding luminal amoebicide, with Metronidazole, the potent tissue amoebicide.

Each film coated tablet of GONAPAN-DF contains  
Diloxanide Furoate B P 500mg  
Metronidazole B P 400mg  
Reko's specs

Each 10ml of GONAPAN-DF suspension contains  
Diloxanide Furoate B P 250mg  
Metronidazole Benzoate B P 320mg  
Equivalent to  
Metronidazole B P 200mg  
Reko's specs

## INDICATIONS

- Acute Amoebiasis
- Chronic Amoebiasis
- Hepatic Amoebiasis and other systemic diseases due to *Entamoeba histolytica*
- Giardiasis

## DOSAGE

### GONAPAN-DF TABLET

Recommended dose for adults is one tablet thrice daily for 5 days, if desired by the physician. **GONAPAN-DF** can be given for 10 days in refractory cases.

### GONAPAN-DF SUSPENSION

The recommended dosage for children aged one to five years

5ml (one teaspoonful) thrice daily for 5 days

### CHILDREN AGED 5-12 YEARS:

5-10ml (1-2 teaspoonful) thrice daily for 5 days

### 12 YEARS AND ABOVE:

10-20ml (2-4 teaspoonful) thrice daily for 5 days

In refractory cases the treatment can be continued up to 10 days, if desired by the physician. Prolonged administration e.g. as a prophylaxis is not recommended.

## CONTRA-INDICATIONS & WARNINGS

There are no absolute contra-indications for the use of Gonapan-DF. It is well tolerated but flatulence and mild gastrointestinal disturbances may occur. Avoid during pregnancy. Alcohol intolerance has been observed in many patients taking Metronidazole. The anticoagulant effect of warfarin may be enhanced by Metronidazole.

## TREATMENT OF OVER DOSAGE:

In case of over dosage, early gastric lavage is recommended.

## PRESENTATION

Gonapan-DF Tablet 3x5's Blister packed in a carton

Gonapan-DF Suspension 60ml bottle



Manufactured by:

**Reko Pharmacal (Pvt) Ltd.**

13-km Multan Road, Lahore, Pakistan



REKO